

Vortex avalanches and the onset of superfluid turbulence

N. B. Kopnin

Low Temperature Laboratory, Helsinki University of Technology, P.O. Box 2200, FIN-02015 HUT, Finland,
L. D. Landau Institute for Theoretical Physics, 117940 Moscow, Russia

(Dated: February 2, 2008)

Quantized circulation, absence of Galilean invariance due to a clamped normal component, and the vortex mutual friction are the major factors that make superfluid turbulence behave in a way different from that in classical fluids. The model is developed for the onset of superfluid turbulence that describes the initial avalanche-like multiplication of vortices into a turbulent vortex tangle.

PACS numbers: 67.57.Fg, 67.40.Vs, 47.37.+q, 47.37.+q, 47.32.Cc

Turbulence in classical liquids [1] is governed by the interplay of the inertial (the first) and viscous (the second) terms in the r.h.s. of the Navier-Stokes equation:

$$\frac{\partial \mathbf{v}}{\partial t} + \nabla \tilde{\mu} = \mathbf{v} \times \boldsymbol{\omega} + \nu \nabla^2 \mathbf{v}, \quad (1)$$

where $\boldsymbol{\omega} = \nabla \times \mathbf{v}$, $\tilde{\mu} = \mu + v^2/2$, and μ is the chemical potential in the case of an isothermal flow. The transition to turbulence is determined by the Reynolds number $Re = RU/\nu$ formed by the characteristic velocity U of the flow and the characteristic size R of the system together with the kinematic viscosity ν . For small Reynolds numbers, the viscous term $-\nu k^2 \mathbf{v}_{\mathbf{k}}$ for a perturbation with a wave vector $\mathbf{k}$ stabilizes the laminar flow. On the contrary, at large $Re \gg 1$ the effect of the inertial term in Eq. (1) is dominating, and laminar flow becomes increasingly unstable towards formation of a chaotic flow of eddies. The evolution of turbulence is described by the Kolmogorov energy cascade: the kinetic energy of the flow is transferred to smaller and smaller length scales, via decay into smaller vortex loops along the Richardson cascade [2], until a scale is reached where the energy is dissipated by viscosity.

In superfluids turbulence acquires new features: Firstly, because it consists of two inter-penetrating components, i.e., the frictionless superfluid and the viscous normal fractions, and secondly, because the vorticity of the superfluid component is quantized in terms of the elementary circulation quantum $\kappa = 2\pi\hbar/M$ where M is a mass of a “superfluid particle” (for ^3He , it is the mass of a pair of atoms, $M = 2m$). If both the normal and the superfluid components are moving, the turbulent state bears more resemblance to the turbulence of classical viscous liquids. The superfluid turbulence is now a subject of intensive studies, see [3] for a review. A new class of turbulent flow becomes possible when the normal component is so viscous that it is essentially immobile while the main flow is supported by the superfluid component which contains a large number of quantized vortices. We refer to this state as *one-component superfluid turbulence*. A superfluid motion of such type can be realized for flows in narrow channels and in rotating

superfluids. The velocity of superflow can be characterized by the “superfluid Reynolds number” $Re_s = U_s R/\kappa$ where U_s is the mean superfluid velocity with respect to the normal component (counterflow velocity). The condition $Re_s \sim 1$ corresponds to the Feynman criterion at which it becomes energetically favorable to unpin a vortex from the container wall. If a large nucleation barrier for vortex nucleation exists, then vortices are not necessarily created even at high superfluid Reynolds numbers $Re_s > 1$ and the superfluid remains in a vortex-free counterflow state. For velocities well below the intrinsic critical velocity [4] $v_c \sim \kappa/\xi$, superfluid turbulence can be initiated if quantized vortices are injected by some extrinsic means into the flow. The turbulence develops when initial vortices start to multiply and form a vortex tangle. Recent experiments in rotating $^3\text{He-B}$ [5] have revealed qualitatively new conditions for the onset of one-component superfluid turbulence: In contrast to the normal fluids, the superfluid turbulence develops below certain temperature independent of the rotation velocity provided the latter is sufficiently high. According to the measurements of Ref. [5], a few seed vortices injected into the $^3\text{He B}$ counterflow triggered a transition into a state with a turbulent vortex tangle for temperatures $T < 0.6T_c$ while the same injection at $T > 0.6T_c$ did not create any substantial number of vortices in the final state. In these measurements, Re_s as high as 200 have been reached without any noticeable dependence of the transition temperature on the initial counterflow velocity. The theoretical arguments supported by numerical simulations explain [5] these conditions in terms of the mutual friction between the normal and superfluid components that appears in the presence of quantized vortices.

In this Letter we develop a model that describes the onset of one-component superfluid turbulence employing the ideas put forward in Ref. [5]. We consider here the initial multiplication of vortex loops immediately after injection of seed loops into a superflow with $Re_s \gg 1$. Since superfluid vorticity is quantized, formation of vortices during the onset of turbulence is the key issue. We consider this process taking into account effects of mutual friction and derive an equation for evolution of the vortex loop density at the initial stage of the transition

to turbulence.

Quantized vortices mediate transfer of momentum between the normal and superfluid components which produces the mutual friction force on a unit volume of superfluid [6, 7]

$$\mathbf{f}_{\text{mf}} = -\alpha' \rho_s [\mathbf{v}_s \times \boldsymbol{\omega}_s] + \alpha \rho_s [\hat{\boldsymbol{\omega}}_s \times [\boldsymbol{\omega}_s \times \mathbf{v}_s]] \quad (2)$$

for a normal component stationary in the container frame, $\mathbf{v}_n = 0$. Here $\boldsymbol{\omega}_s = \nabla \times \mathbf{v}_s$ and $\hat{\boldsymbol{\omega}}_s$ is the unit vector. For an array of quantized vortices

$$\boldsymbol{\omega}_s = \kappa \sum_n \int d\mathbf{r}_n \delta(\mathbf{r} - \mathbf{r}_n)$$

where $\mathbf{r}_n$ is the coordinate of the n -th vortex line, and $\delta(\mathbf{r} - \mathbf{r}_n)$ is a three-dimensional δ -function. The dimensionless parameters α and α' describe the mutual friction between the superfluid and normal components. The mutual friction typically arises from interaction between quasi-particle excitations and the vortices [7, 8]. The energy dissipation is determined by the viscous mutual friction coefficient α in Eq. (2), and α' is the reactive coefficient. Including the mutual friction force in the Euler equation for superfluid velocity we arrive at [7]

$$\frac{\partial \mathbf{v}_s}{\partial t} + \nabla \tilde{\mu} = (1 - \alpha') \mathbf{v}_s \times \boldsymbol{\omega}_s + \alpha \hat{\boldsymbol{\omega}}_s \times (\boldsymbol{\omega}_s \times \mathbf{v}_s) \quad (3)$$

where $\alpha, 1 - \alpha' > 0$.

Regarding the onset of turbulence, the reactive coefficient α' in Eq. (3) simply renormalizes the inertial term of conventional hydrodynamics [the first term in r.h.s. of Eq. (1)]. Indeed, performing the stability analysis of Eq. (3), we would write $\mathbf{v}_s = \mathbf{v}_{s0} + \mathbf{v}'_s$, etc., where $\mathbf{v}_{s0}$ is the solution of a stationary problem while $\mathbf{v}'_s(t)$ is a time-dependent perturbation and obtain a first-order in time linear differential equation for $\mathbf{v}'_s(t)$ with time-independent coefficients made out of $\mathbf{v}_{s0}$ and its spatial derivatives. Making re-scaling of time $t = t'/(1 - \alpha')$ we arrive at the linearized version of Eq. (3) where the first term has now the prefactor unity like the one in the Euler equation [i.e., in Eq. (1) without the viscous term]. The coefficient α in front of the dissipative term [the second term in the r.h.s. of Eq. (3)] is replaced with $q = \alpha/(1 - \alpha') > 0$. Therefore, one concludes that the first term in the r.h.s. of Eq. (3) drives the flow instability towards turbulence in the same way as the inertial term in the Euler equation does for a potential flow in classical hydrodynamics [1]. The fundamental difference from classical turbulence is that the dissipative term in Eq. (3) that stabilizes the flow has now the same scaling dependence on velocity and its gradients as the inertial term. The relative importance of dissipation in formation of turbulence is thus described by the intrinsic dimensionless parameter of the superfluid, $q = \alpha/(1 - \alpha')$ independent of extrinsic quantities such as U_s or R .

The instability leading to multiplication of seed vortices can be derived from Eq. (3). Taking curl of the both sides of Eq. (3) we obtain the vorticity equation

$$\frac{\partial \boldsymbol{\omega}_s}{\partial t} = (1 - \alpha') \nabla \times [\mathbf{v}_s \times \boldsymbol{\omega}_s] + \alpha \nabla \times [\hat{\boldsymbol{\omega}}_s \times [\boldsymbol{\omega}_s \times \mathbf{v}_s]] \quad (4)$$

Let ℓ be a characteristic size of vortex loops. In an entangled vortex-loop state, their three-dimensional density is $n \sim \ell^{-3}$ while the vortex-loop length per unit volume [9] (two-dimensional vortex density) is $L = \ell n = \ell^{-2} = n^{2/3}$. Let us express the two terms in the r.h.s. of Eq. (4) through the vortex density L . Since $\omega_s \sim \kappa L$, the first term in the r.h.s. that drives the instability becomes

$$\dot{L}_+ \sim (1 - \alpha') v_s L^{3/2} \sim (1 - \alpha') (U_s - v_0) L^{3/2} \quad (5)$$

The superfluid velocity in Eq. (5) is assumed to be $v_s = U_s - v_0$ where U_s is the counterflow velocity, and $v_0 \sim \kappa/\ell$ is the self-induced velocity for a vortex loop of a length ℓ . The kinetic energy of superfluid grows due to the increase in the loop density. The kinetic energy is taken from the external source at the length scale R and is transferred to smaller scales. The energy flow into the unit volume of the fluid is $dE/dt \sim E_L \dot{L}_+$ where E_L is the energy of the vortex per unit length. However, the energy transfer goes on only during the transient period when the vortex-loop density increases. When the loop density reaches a value such that $U_s = v_0 \sim \kappa L^{1/2}$ the vortex multiplication saturates and the energy transfer stops. If the density happens to become larger, it will decrease towards $L_{\text{sat}} \sim (U_s/\kappa)^2$ while the kinetic energy is returned from smaller to larger scales and, finally, back to the external source. In other words, saturation is reached when the “turbulent superfluid Reynolds number” $\text{Re}_s^{(\text{turb})} = U_s \ell / \kappa$ is of the order of unity. The condition $\text{Re}_s \gg 1$ ensures the separation of scales $\ell \ll R$ that is required for formation of a vortex tangle. The limit of large $\text{Re}_s \propto \kappa^{-1}$ is equivalent to vanishing Planck constant since $\kappa \propto \hbar$ so that the vorticity becomes a continuous variable like in classical fluids. However, the saturation takes place only in quantum superfluids: L_{sat} diverges for classical fluids where κ can vanish. This difference can be seen in the behavior of the turbulent velocity in normal fluids [1]: the velocity at a length scale λ decreases as $v_\lambda \propto \lambda^{1/3}$ according to the Kolmogorov scaling law. On the contrary, the velocity in a superfluid turbulence scales as $v_\ell \sim \kappa/\ell$ for smaller ℓ . This shows that the “effective circulation” in normal fluid decreases, $\kappa_{\text{eff}} \propto \lambda^{4/3}$. The lower limit on λ in classical fluids is determined by viscosity at the Kolmogorov dissipation scale $\lambda_0 \sim R/\text{Re}^{1/4}$ [1].

The multiplication of vortex loops described by Eq. (5) can be understood in terms of vortex collisions and interconnections. Such processes were indeed seen in numerical simulations on quantized vortices [5, 10, 11]. Reconnection of vortices accompanied by formation of vortex

tangle in normal fluids were considered recently in Refs. [12, 13]. Each reconnection of quantized vortices takes a microscopic time of the order of the quasiparticle collision time, much shorter than the times involved in hydrodynamic processes. It is accompanied by a small dissipation within a volume ξ^3 where ξ is the core radius of the order of the superfluid coherence length. We consider these processes as instantaneous and neglect the corresponding dissipation. The rate of increase in the vortex loop density should be quadratic in n thus $\dot{n}_+ = Av_r n^2 \ell^2$. Here v_r is the relative velocity of the vortex loops, ℓ^2 is the loop cross section, a constant $A \sim 1$ describes an “efficiency” of the vortex multiplication due to pair collisions. Using definition of L the vortex multiplication rate becomes $\dot{L}_+ \sim v_r L^{3/2}$. The vortex velocity is determined through the mutual friction parameters α and α' such that $\mathbf{v}_L = (1 - \alpha')\mathbf{v}_s - \alpha\hat{\omega}_s \times \mathbf{v}_s$. The relative velocity of loops v_r is proportional to the longitudinal component of $\mathbf{v}_L$: $v_r \sim (1 - \alpha')v_s$ which agrees with Eq. (5).

We assume that vortex multiplication occurs first within a certain region in the fluid near the location of the seed loops from where the created vortex tangle penetrates into the rest volume of the fluid. The effect of the second (viscous) term in the r.h.s. of Eq. (4) is to decrease the loop density in the multiplication region by inflating the loops due to counterflow and extracting them from this region. The viscous component of the mutual friction force leads to variation in the vortex loop length $\dot{\ell} \sim 2\pi v_L \sim \alpha(U_s - v_0)$. The length increases while the density decreases if $U_s - v_0 > 0$. When $v_0 > U_s$, the density increases since the loops shrink due to the friction. Thus the rate of vortex-loop density variation due to the viscous component becomes

$$\dot{L}_- \sim -\alpha(U_s - v_0)L^{3/2}. \quad (6)$$

The loops expanding according to Eq. (6) are extracted from the region of their multiplication into the bulk if $U_s > v_0$ or vice versa if $U_s < v_0$. This process is accompanied by dissipation of the kinetic energy. The dissipation persists until the equilibrium density is reached.

As we see, both inertial and viscous mutual friction terms, Eqs. (5) and (6), have the same dependence on the vortex density, i.e., on the vortex length scale. The total variation of the loop density in the multiplication region is the sum of the two processes, $\dot{L} = \dot{L}_+ + \dot{L}_-$. Putting $v_0 = \kappa/\ell = \kappa L^{1/2}$ we obtain

$$\dot{L} = \beta \left[U_s L^{3/2} - \kappa L^2 \right] \quad (7)$$

where $\beta = A(1 - \alpha') - \alpha = (A - q)/(1 - \alpha')$.

Equation (7) looks like the Vinen equation [9] for superfluid turbulence. However, the difference is that the coefficient β can now have either positive or negative sign depending on the mutual friction parameters. One distinguishes two limits. In the viscosity dominated regime

when $\beta < 0$, the rate of extraction of vortex loops exceeds the rate of multiplication; there is no time for vortices to multiply since all the seed and newly created vortices are immediately wiped away into the bulk fluid. The number of vortices in the final state is essentially equal to the number of initial vortices, and the turbulent state is not formed. The corresponding stable solution to Eq. (7) is $L \rightarrow 0$. Equation (7) has another point of attraction, $L \rightarrow \infty$. This would correspond to a decay of initially created vortex tangle in a situation without net counterflow: the vortex loops shrink due to the viscous mutual friction force.

In the low-viscosity regime when $\beta > 0$, the rate of multiplication is faster, and the number of created vortex loops is large: Each new vortex loop serves as a source for producing more vortices. As a result, an avalanche-like multiplication takes place, which leads to formation of a turbulent vortex tangle. As the number of vortex loop grows, the self-induced velocity increases and finally the saturated density L_{sat} is reached. Indeed, Eq. (7) has the stable solution $L_{sat} = (U_s/\kappa)^2$ which corresponds to the Vinen equilibrium vortex density for the counterflow turbulence [14]. For $\beta > 0$, the solution to an equation of the type of Eq. (7) for a constant U_s was found in Ref. [14]. At the initial stage, when $L \ll L_{sat}$, the vortex density in the multiplication region varies as

$$L/L_{sat} = \left[(L_{sat}/L_0)^{1/2} - (t/\tau) \right]^{-2} \quad (8)$$

with the rate

$$\tau^{-1} = \beta U_s^2 / 2\kappa.$$

In Eq. (8), L_0 is the (small) initial vortex density at $t = 0$. The time dependence of L turns to exponential $L_{sat} - L \propto \exp(-t/\tau)$ for $L \rightarrow L_{sat}$. The time τ decreases with lowering the temperature since β increases. The energy $E_L \dot{L}_+$ pumped into the fluid goes first to a reversible increase in the kinetic energy of the superflow. A part of it is then dissipated due to the mutual friction.

The overall evolution of the vortex density can be seen as an interplay of two processes. The first is the turbulent instability governed by Eq. (5). The second process is expansion of vortex loops from the multiplication region into the bulk due to the viscous part of the mutual friction force. Equation (6) taken with the *opposite* sign gives the rate of vortex-loop-density flow *into the bulk*. In this form, it coincides exactly with the Vinen equation [9] as derived by Schwarz [10]. The solution for a constant U_s has the form of Eq. (8) with the characteristic rate

$$\tau_b^{-1} = \alpha U_s^2 / 2\kappa.$$

This time increases with lowering the temperature since α decreases.

The instability develops out of initial seed vortices introduced into the fluid. In experiments [5] these seed vortices were produced by the Kelvin-Helmholtz instability

at the interface between the A and B phases in rotation [15]. We note that the inequality $q \ll 1$ is also the condition for propagation of the Kelvin waves [16, 17]. Perturbations of the seed vortex lines in the form of Kelvin waves can grow and multiply under the action of longitudinal flows. These processes obviously belong to the same class and can also be described within our model.

Consider a container of a radius R rotating with an angular velocity Ω . If the counterflow velocity were kept constant $U_s = \Omega R$ as in the vortex-free state, the density L_{sat} would correspond to the number of vortices in the container $N_{max} \sim \pi R^2 L_{sat} \sim N_{eq}^2$ where $N_{eq} = 2\pi R^2 \Omega / \kappa$ is the equilibrium number of vortices. Such a big number N_{max} of vortices is unstable, therefore it relaxes towards N_{eq} . This decay of turbulence takes much longer times than τ and is accompanied by polarization of vortex loops and screening of the counterflow, which is beyond the scope of the present paper. In the final state, with the vortex cluster formed in the center of the container, the counterflow velocity is decreased down to $U_s \sim \kappa / r_0 \sim (\kappa \Omega)^{1/2}$ where r_0 is the intervortex distance for the equilibrium density $L_{eq} \sim \Omega / \kappa$.

In Fermi superfluids and superconductors, the parameter $q \approx (\omega_0 \tau)^{-1}$, where ω_0 is the spacing between the bound states of quasiparticles in the vortex core and τ^{-1} is their lifetime broadening due to scattering from the normal component [8]. The ratio $\alpha / (1 - \alpha')$ decreases exponentially [8, 18, 19] at low temperatures. As a result, below a certain temperature, an avalanche-like burst of vortices in the multiplication region up to the density L_{sat} can be achieved, and the well-developed turbulence can be realized. However, a large number of vortices cannot be created for higher temperatures. In $^3\text{He-B}$, q varies from zero, at $T \rightarrow 0$, to ∞ , at $T \rightarrow T_c$, with $q \sim 1$ at $T \sim 0.6T_c$ [19]. As pointed out in Ref. [5], this temperature is in a good agreement with the experimental observations of the onset of turbulence. In superfluid He-II, the friction coefficient is always small except for the immediate vicinity of T_λ . Thus, an entangled turbulent vortex state is usually formed. On the contrary, in $^3\text{He-A}$ the friction parameter α is large except for low temperatures [18] so that the turbulent state would be possible only for very low temperatures.

To summarize, we develop a model for the onset of superfluid turbulence taking into account the mutual friction between the normal and superfluid components that appears in the presence of quantized vortices. The condition for turbulent instability is velocity independent: If the viscous component of the mutual friction force is small, which happens at low temperatures, the avalanche-

like multiplication of seed vortex loops takes place. The created vortex loops form a dense vortex tangle which further evolves into a developed turbulent flow. On the contrary, the number of created vortex loops is low and a turbulent state is not formed in the viscosity dominated regime realized at higher temperatures.

I am grateful to V. Eltsov, M. Krusius, and G. Volovik for many stimulating discussions. This work was supported by Russian Foundation for Basic Research.

-
- [1] L.D. Landau and E.M. Lifschitz, *Fluid mechanics*, Pergamon, Oxford, 1987.
 - [2] U. Frisch *Turbulence: The legacy of A.N. Kolmogorov*, Cambridge University Press, 1995.
 - [3] W.F. Vinen, J.J. Niemela, J. Low Temp. Phys. **128**, 167 (2002).
 - [4] V.M.H. Ruutu, Ü. Parts, J.H. Koivuniemi, N.B. Kopnin, and M. Krusius, J. Low Temp. Phys. **107**, 93 (1997).
 - [5] A. P. Finne, T. Araki, R. Blaauwgeers, V. B. Eltsov, N. B. Kopnin, M. Krusius, L. Skrbek, M. Tsubota, and G. E. Volovik, Nature, **424**, 1022 (2003).
 - [6] H.E. Hall and W.F. Vinen, Proc. Roy. Soc. A **238**, 215 (1956).
 - [7] E.B. Sonin, Rev. Mod. Phys. **59**, 87 (1987).
 - [8] N. B. Kopnin, Rep. Progr. Phys. **65**, 1633 (2002).
 - [9] W. F. Vinen, Proc. R. Soc. London, Ser. A **242**, 493 (1957).
 - [10] K.W. Schwarz, Phys. Rev. B **38**, 2398 (1988).
 - [11] M. Tsubota, T. Araki, and C.F. Barenghi, Phys. Rev. Lett. **90**, 205301 (2003).
 - [12] D. Kivotides and A. Leonard, Phys. Rev. Lett. **90**, 234503 (2003).
 - [13] P. Chatelain, D. Kivotides, and A. Leonard, Phys. Rev. Lett. **90**, 054501 (2003).
 - [14] K.W. Schwarz and J.R. Rozen, Phys. Rev. B **44**, 7563 (1991).
 - [15] R. Blaauwgeers, V. B. Eltsov, G. Eska, A. P. Finne, R. P. Haley, M. Krusius, J. J. Ruohio, L. Skrbek, and G. E. Volovik, Phys. Rev. Lett. **89**, 155301 (2002).
 - [16] C. F. Barenghi, R. J. Donnelly, and W. F. Vinen, Phys. Fluids **28**, 498 (1985).
 - [17] W.F. Vinen, M. Tsubota, and A. Mitani, Phys. Rev. Lett. **91**, 135301 (2003).
 - [18] N. B. Kopnin and A. V. Lopatin, Phys. Rev. B **56**, 766 (1997).
 - [19] T. D. C. Bevan, A. J. Manninen, J. B. Cook, A. J. Armstrong, J. R. Hook, and H. E. Hall, Phys. Rev. Lett. **74**, 750 (1995); **74**, 3092 (1995); T. D. C. Bevan, A. J. Manninen, J. B. Cook, H. Alles, J. R. Hook, and H. E. Hall, Journ. Low Temp. Phys. **109**, 423 (1997).